

## References and Recommended Readings:

Brady D, Sun LH. 2014. FDA found more than smallpox in storage room. Washington Post (Washington, DC) 17 July: A1.

Finkelstein MM, Verma DK, 2001. Exposure estimation in the presence of nondetectable values: another look. *AIHAJ*. Mar-Apr; 62 (2):195-8

Helsel DR, 2005. More than obvious: better methods for interpreting nondetect data. *Environ Sci Technol*. Oct 15;39 (20): 419A-423A.

Hornung WR, Reed LD, 1980. Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg* 5:46-51

Kim M.Y, Dubin N, 1995. Study Design and Sample size Considerations for Half-Life Studies. *Arch. Environ. Contam. Toxicol*. 30, 423-429

Lubin JH, Colt JS, Camann D, Davis S, Cerhan JR, Severson RK, Bernstein L, Hartge P, 2004. Epidemiology Evaluation of Measurements Data in the Presence of Detection Limits. *Environmental Health Perspectives*. Vol. 112 number 17

Lynch JR, Ayer HE. 1966. Measurement of Dust Exposures in the Asbestos Textile Industry. *Amer. Industrial Hyg. Assoc. Journal*. Sep – Oct; 27 (5): 431 – 7.

National Academy of Science, Committee on Human Biomonitoring for Environmental Toxicants, National Research Council. *Human Biomonitoring for Environmental Chemicals*. (2006) ISBN: 0-309-66315-6, 316 pages, 6x9.

NCI Best Practices for Biospecimen Resources.

<http://biospecimens.cancer.gov/bestpractices/2011-NCIBestPractices.pdf>

Rothman N, Stewart WF, Schulte PA. 1995. Incorporating Biomarkers into Cancer Epidemiology: A Matrix of Biomarker and Study Design Categories. *Cancer Epidemiology Biomarkers & Prevention*; AACR journal. Jun;4 (4): 301 – 11.